

The surjectivity problem for one-generator, one-relator extensions of torsion-free groups

MARSHALL M COHEN
COLIN ROURKE

Cornell University, Ithaca NY, 14853-4102, USA
Warwick University, Coventry, CV4 7AL, UK

Email: marshall@math.cornell.edu and cpr@maths.warwick.ac.uk

Abstract We prove that the natural map $G \rightarrow \widehat{G}$, where G is a torsion-free group and \widehat{G} is obtained by adding a new generator t and a new relator w , is surjective only if w is conjugate to gt where $g \in G$. This solves a special case of the surjectivity problem for group extensions, raised by Cohen [2].

AMS Classification 20E22, 20F05; 57M20, 57Q10

Keywords Surjectivity problem, torsion-free groups, Whitehead torsion, Kervaire conjecture

1 Introduction

In this paper we prove the following theorem.

Main Theorem *Suppose that G is a torsion-free group and that $\langle t \rangle$ is an infinite cyclic group with generator t . Let w be an element of the free product $G * \langle t \rangle$ and let $\langle\langle w \rangle\rangle$ be the normal subgroup of $G * \langle t \rangle$ generated by w . View G as a subgroup of $G * \langle t \rangle$ and let i be the inclusion $G \rightarrow G * \langle t \rangle$. Consider the natural homomorphism*

$$q = \pi i: G \xrightarrow{i} G * \langle t \rangle \xrightarrow{\pi} \widehat{G} = \frac{G * \langle t \rangle}{\langle\langle w \rangle\rangle}.$$

If q is onto then w is conjugate to gt for some $g \in G$.

There are standard ways in which this algebraic situation may be realized topologically. These lead to the following results.

Corollary 1 *Suppose that L is a connected CW complex with torsion-free fundamental group and that the CW complex $\widehat{L} = L \cup e^1 \cup e^2$ is constructed by attaching a 1-cell to L and a 2-cell to $L \cup e^1$. If the inclusion map $j: L \rightarrow \widehat{L}$ induces a surjection $j_*: \pi_1 L \rightarrow \pi_1 \widehat{L}$ then j is a simple-homotopy equivalence.*

Proof This follows from elementary facts about the invariance of Whitehead torsion under homotopy of attaching maps [1; Section 5] and the fact that if $\widehat{L} = L \cup e^1 \cup e^2$, where e^1 is a circle and e^2 is a 2-cell attached by a word gt , then e^1 is a free face of \widehat{L} and \widehat{L} collapses to L by an elementary collapse. \square

Corollary 2 *Suppose that M is a connected n -manifold with torsion free fundamental group and that (W, M, M') is an h -cobordism with exactly one handle of index one and one handle of index two and no other handles (or dually with exactly one n -handle and one $(n - 1)$ -handle). Then (W, M, M') is an s -cobordism.*

Proof This is a consequence of the fact that a k -handle $D^k \times D^{n+1-k}$ collapses to its core union its attaching tube, $D^k \times \{0\} \cup \partial D^k \times D^{n+1-k}$, see eg [10; Chapter 6]. So the CW theory applies to the handlebody theory. \square

Background

The surjectivity problem for group extensions and the question of which Whitehead torsions can be realized were formulated by Cohen [2] and Metzler [6]; for more details on these problems and the relevance of our results, see Section 5.

It will be useful to note from the outset that the conclusion of the main theorem may be restated according to the following lemma.

Lemma 1 *If $G \subset G * \langle t \rangle \xrightarrow{\pi} G * \langle t \rangle / \langle\langle W \rangle\rangle$ where W is a set of words in $G * \langle t \rangle$ then $q = \pi|_G$ is onto $\iff gt$ lies in the kernel of π for some $g \in G$.*

Proof q is onto $\iff [\pi(t) \in \pi(G)] \iff [\pi(t) = \pi(g^{-1}) \text{ for some } g \in G] \iff [\pi(gt) = 1] \iff [gt \in \text{kernel}(\pi) \text{ for some } g \in G].$ \square

Any element $w \in G * \langle t \rangle$ has a unique expression as a *reduced word*, $w = g_0 t^{q_1} g_1 t^{q_2} \dots g_{n-1} t^{q_n} g_n$, where $g_i \in G$ are non-trivial for $0 < i < n$ and q_i are non-zero integers for each i . The word w is *cyclically reduced* if further $g_n = 1$ and if $n > 1$ then $g_0 \neq 1$. Up to cyclic permutation there is a unique cyclically reduced word in the conjugacy class of w , see eg [3, Proposition 3.9]. Since \widehat{G} depends only the conjugacy class of w , there is no loss in assuming that w is cyclically reduced and we shall do so without comment from now on. We call $\sum_{i=1}^n q_i$ the *exponent sum* of t in w , denoted $\text{ex}(w)$. The unreduced word $t^{q_1} t^{q_2} \dots t^{q_n}$ is called the *t -shape* of w and, thinking of $w = 1$ as an equation over G , we call the elements g_i the *coefficients* of w .

It is easy to see that if $q: G \longrightarrow \widehat{G}$ is surjective then $\text{ex}(w) = \pm 1$, since otherwise the abelianization of $\widehat{G}/(q(G) = 1)$ will be non-trivial. So, replacing w by w^{-1} if necessary, we may assume in our discussion that $\text{ex}(w) = 1$. Under this hypothesis Klyachko [8], in 1993, gave a brilliant argument to prove the following theorem, which implies the Kervaire conjecture [7] in the case where G is torsion-free.

Theorem (Klyachko) *If G is a torsion-free group and $w \in G * \langle t \rangle$ with $\text{ex}(w) = 1$ then the natural homomorphism $q: G \longrightarrow \widehat{G} = \frac{G * \langle t \rangle}{\langle\langle w \rangle\rangle}$ is injective.*

An exposition (and extension) of Klyachko's theorem was given by Fenn and Rourke [4] in 1996. To prove our theorem we will use Klyachko's result and his method, following closely the exposition in [4]. We will quote some definitions and results from [4] and give those proofs in detail for which the arguments differ and for which (proving the contrapositive) the hypothesis is used that w is not conjugate to gt for any $g \in G$.

Outline of the paper

In Section 2 we consider a group Γ in a slightly more general situation than G above. We assume (contrary to our Main Theorem) that w is not conjugate to gt for any $g \in \Gamma$ but that some gt is in the kernel of $\Gamma * \langle t \rangle \longrightarrow \widehat{\Gamma}$. We show how to construct a certain non-trivial CW subdivision of the 2-sphere, with edges labelled $t^{\pm 1}$ and all but one corner labelled by an element of Γ .

In Section 3 we prove our main theorem in a special case: We denote $g^t = t^{-1}gt$. If w has the form $w = b_0 a_0^t b_1 a_1^t \dots b_r a_r^t c t$, where the a_i, b_i and c are all elements of G and $b_0 a_0^t b_1 a_1^t \dots b_r a_r^t c \notin G$ then $q: G \longrightarrow \widehat{G}$ is not onto.

In Section 4, we complete the proof of the main theorem. We use an algebraic trick to parlay the result of Section 4 into a proof that, in general, if $\text{ex}(w) = 1$ and w is not conjugate to gt for any $g \in G$ then $q: G \longrightarrow \widehat{G}$ is not onto.

In Section 5 we briefly discuss the general surjectivity problem, in which n generators and n relators are added to a group G . We give a bit of history and comment on the relevance of our result for $n = 1$ to the general problem.

Finally, in Section 6 we give an extension of our result and ask a question.

2 The cell subdivision lemma

In this section we prove the cell subdivision lemma (below) which is modelled on [4; Lemma 3.2].

The lemma uses the idea of a *corner* of a 2-cell in a cell subdivision K of the 2-sphere. This can be regarded as the (oriented) angle formed by the two adjacent edges meeting at a vertex (0-cell) in the boundary of the 2-cell. If all the corners of a 2-cell are labelled by elements of a group, then a word can be read around the 2-cell boundary by composing these elements either unchanged or inverted according as the orientation of the corner agrees or disagrees with that of the 2-cell boundary. Similarly if all the corners at a vertex are labelled then a word can be read around that vertex. We shall always orient corners *clockwise*, thus if the above words are read *clockwise* for vertices and *anticlockwise* for 2-cells, then no inversion is necessary. See figure 1 for an example: the word read around the boundary is abc ; after insertion of t or t^{-1} at the arrows (see part (e) of the lemma below) it reads $tat^{-1}bt^{-1}c$.




Figure 1: Reading the boundary of a 2-cell: $tat^{-1}bt^{-1}c$

Let H be a subgroup of a group Γ and let $g \in \Gamma$. We say that g is *free relative to H* if the subgroup $\langle g, H \rangle$ of Γ generated by g and H is naturally the free product $\langle g \rangle * H$ of an infinite cyclic group $\langle g \rangle$ with H . (Note in particular that g has infinite order.)

If g, h are elements of a group let g^h denote $h^{-1}gh$.

In this section and the next, we shall consider the following working hypotheses:

Working hypotheses

Suppose that H and H' are two isomorphic subgroups of a group Γ under the isomorphism $h \rightarrow h^\phi$, $h \in H$. Suppose that for each i , a_i, b_i are elements of Γ such that a_i is free relative to H and b_i is free relative to H' . Let c be an arbitrary element of Γ .

Let w_0 be the word

$$b_0 a_0^t b_1 a_1^t b_2 a_2^t \cdots b_r a_r^t c t$$

in $\Gamma * \langle t \rangle$, where $r \geq 0$, and let $W \subset \Gamma * \langle t \rangle$ be the set of words $\{w_0, h^t (h^\phi)^{-1} \mid h \in H\}$. Let $\langle\langle W \rangle\rangle$ be the normal closure of W in $\Gamma * \langle t \rangle$ and let $\widehat{\Gamma}$ denote $(\Gamma * \langle t \rangle) / \langle\langle W \rangle\rangle$.

Cell subdivision lemma

Assume the working hypotheses, above. Suppose that, for some $g \in \Gamma$, gt is in the kernel of the natural map $\Gamma * \langle t \rangle \rightarrow \widehat{\Gamma}$.

Then there is a cell subdivision K of the 2-sphere such that

- (a) the edges (1-cells) of K are oriented,
- (b) the corners (all oriented clockwise) are labelled by coefficients of elements w or w^{-1} for $w \in W$, with the exception of one particular corner at one particular vertex v_0 which is unlabelled,
- (c) the clockwise product of the corner labelling around any vertex is $1 \in \Gamma$ except for v_0 where it is undefined,
- (d) there is one special 2-cell e_∞^2 which contains the unlabelled corner and has boundary a single edge and the single vertex v_0 ,
- (e) with the exception of e_∞^2 , the corner labels of any 2-cell (in anticlockwise order) are the coefficients of w or w^{-1} for some $w \in W$ (up to cyclic rotation) with the property that, if on passing from one corner to an adjacent corner the element t or t^{-1} is inserted according to whether the intervening edge is oriented in the same or opposite direction (see figure 1), then the whole of w or w^{-1} is recovered,
- (f) the cell decomposition is irreducible in the following senses:
 type (1) there do not exist two 2-cells with an edge in common (necessarily read as t in one and t^{-1} in the other) such that, starting with one vertex of this edge, the words read in these 2-cells are inverses of each other,
 type (2) there does not exist a chain of 2-gons with common vertices a, b such that the product of the corner labels in the chain at a (or, equivalently, at b) is $1 \in \Gamma$,
- (g) the cell subdivision is non-degenerate in that there exist at least two vertices and at least three 2-cells; in particular there is a cell $e_1^2 \neq e_\infty^2$ whose boundary contains ∂e_∞^2 as a proper subset (see figure 5).

Proof The proof uses transversality as in the proofs of [4; Lemmas 3.1 and 3.2].

Choose a 2-complex L with $\pi_1(L) = \Gamma$ and form the 2-complex \widehat{L} with $\pi_1(\widehat{L}) = \widehat{\Gamma}$ by attaching a 1-cell γ to the base point $*$ of L (corresponding to t) and a 2-cell σ_w with attaching map determined by w for each $w \in W$.

Since gt is trivial in $\pi_1(\widehat{L})$ there is a map of a 2-disc $f: D^2 \rightarrow \widehat{L}$ whose boundary maps to $L \cup \gamma$ and which represents $gt \in \pi_1(L \cup \gamma) = \Gamma * \langle t \rangle$. Make f transverse to the centres of the 2-cells σ_w . It follows that the inverse images of small neighbourhoods of these centres is a collection of disjoint discs D_1, \dots, D_m in the interior of D^2 . By a radial expansion of f on these discs we may assume that their image is the whole of one of the σ_w . It follows that the punctured disc $P = \text{closure}(D^2 - (D_1 \cup \dots \cup D_m))$ is mapped by f to $L \cup \gamma$. Let p be the centre of γ . Make $f|_P$ transverse to p . Then $f^{-1}p$ is a 1-manifold Z properly embedded in P . By a radial expansion along γ we can assume that Z has a neighbourhood N which is a normal I -bundle, where each fibre is mapped by f to γ and $\text{closure}(P - N)$ is mapped by f to L .

We now simplify the subset $\mathcal{H}_f = D_1 \cup \dots \cup D_m \cup N$ of D^2 as follows. Suppose N contains an annulus component \mathcal{A} in the interior of P . Let D' denote the interior disc of D^2 which bounds the interior boundary component of the annulus. Then $D'' = D' \cup \mathcal{A}$ is a sub disc of D^2 whose boundary is mapped to $*$ by f . We can then redefine f so that $f(D'') = *$ leaving f unchanged outside D'' .

At this point \mathcal{H}_f can be regarded as a collection of 0-handles (the D_i) and 1-handles (the components of N) attached to the 0-handles and to ∂D^2 (in fact there is precisely one 1-handle attached to ∂D^2 by one end) see figure 2.




Figure 2: A view of \mathcal{H}_f

We now prove that we may assume that \mathcal{H}_f is connected. Suppose not. Choose an innermost component C . Draw a simple loop α around C separating it from the rest of \mathcal{H}_f . Up to conjugacy α represents an element of $\pi_1(L) = \Gamma$ which is trivial in $\pi_1(\widehat{L}) = \widehat{\Gamma}$. But Klyachko proves that Γ injects in $\widehat{\Gamma}$ (this is the precise content of [4; Theorem 4.1, page 62]) and hence we may redefine f so that the inside of α is mapped to L , which simplifies \mathcal{H}_f .

Note that the 0–handles can be labelled by elements w or w^{-1} for $w \in W$ according to the corresponding 2–cell of \widehat{L} and orientation. We say that \mathcal{H}_f is *type (1) reducible* if there is a pair of 0–handles labelled by w and w^{-1} (the same w) and joined by a 1–handle which represents the same occurrence of t (respectively t^{-1}) in each word. In this situation we can again simplify \mathcal{H}_f without changing $f|_{\partial D^2}$ by redefining f near these 0–handles and joining 1–handle (see figure 3).




Figure 3: Type (1) reduction of \mathcal{H}_f

We say that \mathcal{H}_f is *type (2) reducible* if there is a chain of 0–handles (each having two 1–handles attached to it) labelled by words $h_i^t (h_i^\phi)^{-1}$, $i = 1, 2, \dots, q$ with $h_1 h_2 \dots h_q = 1$ in Γ (figure 4).




Figure 4: Type (2) reduction of \mathcal{H}_f

If the chain forms a loop, the handlebody is not connected and this chain and everything inside it may be eliminated as indicated earlier. Otherwise, the curve α indicated in figure 4 maps to tt^{-1} in $\Gamma * \langle t \rangle$ and there is another simplification given by omitting this chain of 0–handles and redefining f inside α using the null-homotopy of tt^{-1} in $L \cup \gamma$. After these simplifications there may now be more simplifications of the first two types which can be performed. Repeat all four until no more are possible, Thus we can assume that \mathcal{H}_f is connected and irreducible.

We now extend \mathcal{H}_f to a handle decomposition \mathcal{H} of $S^2 \supset D^2$ by letting the outside of D^2 be one 0–handle (denoted h_∞^2) and the regions of $D^2 - \mathcal{H}_f$ be the 2–handles.

The required 2-complex K is the dual complex to \mathcal{H} obtained by putting a vertex inside each 2-handle and joining by an edge across each 1-handle. The outside 2-cell is e_∞^2 (containing h_∞^2) and has boundary containing a single vertex v_0 . Corners of 2-cells other than this corner are labelled by the coefficient of the word w or w^{-1} labelling the 0-handle inside the 2-cell opposite the corner. See figure 5.




Figure 5: The cell complex K (shown dashed)

The required properties of K all follow from the construction: 1-cells are oriented by the orientations of the I -bundles (1-handles) that they cross and properties (a) to (e) follow at once (the word read around the boundary of a 2-cell is the label on the contained 0-handle). Property (f) follows from the irreducibility of \mathcal{H}_f .

Finally, property (g) uses the hypothesis that $r \geq 0$ (ie, that w_0 is not conjugate to gt for any $g \in \Gamma$). In order that every 1-handle of \mathcal{H} have each end on some ∂D_i^2 , except that one of them has one end on ∂D^2 , at least one of the 0-handles D_i^2 must be a w_0 or w_0^{-1} handle. Because $r \geq 0$, this handle must have at least three 1-handles emanating from it. Thus there have to be at least two 0-handles inside D^2 , so that K has at least three 2-cells. Since the handlebody closes up, $D^2 - \mathcal{H}_f$ must have at least two components, resulting in at least two vertices in K . \square

3 The key technical theorem

In this section we prove the following result whose proof is modeled on that of [4; Theorem 4.1]. We show that the hypotheses of the cell subdivision lemma are self-contradictory.

Key Technical Theorem

*Assume the working hypotheses. Then gt is never in the kernel of the natural map $\Gamma * \langle t \rangle \rightarrow \widehat{\Gamma}$ for any $g \in \Gamma$.*

Remark Assuming this theorem, note that by Lemma 1 in the Introduction, $\Gamma \rightarrow \widehat{\Gamma}$ is not surjective. Therefore by taking H and H' to be trivial, we can now deduce a special case of our main theorem:

If the t -shape of $w = w_0$ is not t (ie, w is not conjugate to gt for any $g \in \Gamma$) but is of the form $t^{-1}tt^{-1} \dots tt^{-1}tt$ then $q: \Gamma \rightarrow \widehat{\Gamma}$ is not surjective.

In the next section we introduce an algebraic trick which will enable us to deduce the general case, where $\text{ex}(w) = 1$ and w is not conjugate to gt for any $g \in G$, from this special case.

Proof The proof relies heavily on the proof and terminology of [4; Theorem 4.1, pages 62–64]. Assume that gt is in the kernel of $\Gamma * \langle t \rangle \rightarrow \widehat{\Gamma}$ where $g \in \Gamma$. By the cell subdivision lemma there is a cell subdivision of S^2 with all 2-cells of the four types I, I', II, II' illustrated in Figure 6 with the exception of the special 2-cell e_∞^2 .

As in [4] we give the two-sphere an orientation (“anticlockwise”) and give each 2-cell of K the induced orientation. A traffic flow is now defined, with a car running around the boundary of each 2-cell in the direction of the induced orientation as follows:

At time 0 let a car on the boundary of a country of type I or I' start at the corner labelled b_0 or b_0^{-1} and proceed in an anticlockwise manner with respect to the orientation of the edge along which it is travelling, moving from corner to corner in unit time except at the corner labelled c or c^{-1} where it stops for $2r - 1$ units. The times when the car is at each corner are illustrated in figure 6. For countries of type II or II' the car starts at the corner labelled h^ϕ or $(h^\phi)^{-1}$ and proceeds in an anticlockwise manner moving from corner to corner in unit time.

For e_∞^2 we need to consider also the 2-cell e_1^2 whose boundary properly contains the boundary of e_∞^2 , see figure 5. Let A be the car on the boundary of e_∞^2 and




Figure 6: The 2-cells I, I', II and II'

B the car on the boundary of e_1^2 . Choose a point ω on ∂e_∞^2 different from v_0 . We shall engineer crashes between A and B to occur precisely at ω . Suppose that B is approaching ω then let A approach ω from the opposite direction to crash at ω . After the crash, let A dawdle near ω until B moves off ∂e_∞^2 (which it must, because ∂e_1^2 properly contains ∂e_∞^2); then let A speed round to just before ω where it again dawdles until B again approaches ω at which point the cycle repeats.

Recall from [4] that a *complete car crash* is said to occur when two cars meet in the interior of an edge (necessarily going in opposite directions) or when a N cars from N neighbouring countries all meet at a vertex of valency N .

Notice that, on ∂e_∞^2 , the given flow has the property that complete crashes occur at ω and nowhere else; in particular, no complete crash occurs at v_0 . However there must be another complete crash occurring at some other vertex of K . (This is for exactly the same reasons as in the proof of [4; Theorem 4.1]. Properties 1 to 4 on pages 63–64 hold here also and the flow satisfies the conditions of the Crash Theorem with Stops [4, Theorem 2.3, page 56]. So there must be another complete crash and, as in [4], this must occur at a vertex.)

This leads to the identical contradiction as on [4; page 64]: The flow has been chosen so that, at a vertex where all the cars come together at the same time, the labels around the corners are all $\{a, a^{-1}\}$ for some coefficient $a = a_i$ of w_0 together with elements of H or $\{b, b^{-1}\}$ for some coefficient $b = b_i$ of

w_0 together with elements of H' . For definiteness assume that we are in the former situation. Then we can read an (unreduced) word of the form $a^\epsilon h_1 h_2 \dots h_{i_1} a^\epsilon h_1 h_2 \dots h_{i_2} a^\epsilon \dots$ which is 1 in Γ . Now if this word contains a subword of the form $a^\epsilon a^{-\epsilon}$ then K is type (1) reducible and if it contains a subword of the form $h_1 h_2 \dots h_i$ which is 1 in Γ then K is type (2) reducible. Since K is irreducible neither of these happen and the word either gives a non-trivial relation in $\langle a, H \rangle$ contradicting the assumption that a is free relative to H or reads $(a^\epsilon)^N = 1$ for $N \geq 1$ which also contradicts the assumption that a is free relative to H (and in particular has infinite order). \square

4 Proof of the main theorem

In the light of the discussion in the Introduction and Lemma 1, we assume that $\text{ex}(w) = 1$ and, proving the contrapositive of the Main Theorem, we assume that w is not conjugate to gt for any $g \in G$. We must prove that

$$gt \text{ is not in the kernel of } \pi: G * \langle t \rangle \longrightarrow \widehat{G} \text{ for any } g \in G. \quad (*)$$

We now use Klyachko's algebraic trick described on pages 64–66 of [4].

Consider the homomorphism $\text{ex}: G * \langle t \rangle \rightarrow \mathbb{Z}$. It is well known that K , the kernel of ex , is a free product of copies of G generated by elements of the form $g^{t^{\mathcal{O}}} = t^{-\mathcal{O}} g t^{\mathcal{O}}$, $1 \neq g \in G$.

Any element of K has a canonical expression of the form $k = g_1^{t^{\mathcal{O}_1}} \dots g_r^{t^{\mathcal{O}_r}}$, where $\mathcal{O}_i \neq \mathcal{O}_{i+1}$ for each i . We shall call the $g_i^{t^{\mathcal{O}_i}}$ the *canonical elements* of k . Let $\min(k)$ be the minimum value of \mathcal{O}_i , $i = 1, \dots, r$ and $\max(k)$ the maximum value. Fix a positive integer m . Consider the following subgroups of K :

$$\begin{aligned} H &= \langle k \in K \mid \min(k) \geq 0, \max(k) \leq m - 2 \rangle \\ H' &= \langle k \in K \mid \min(k) \geq 1, \max(k) \leq m - 1 \rangle \\ J &= \langle k \in K \mid \min(k) \geq 0, \max(k) \leq m - 1 \rangle \end{aligned}$$

and the following subsets:

$$\begin{aligned} X &= \{k \in K \mid \min(k) = 0, \max(k) \leq m - 1\} \\ Y &= \{k \in K \mid \min(k) \geq 0, \max(k) = m - 1\} \\ Z &= \{k \in K \mid \min(k) \geq 1, \max(k) = m\}. \end{aligned}$$

Lemma 2 [4; Lemma 4.2, page 65] *Let $w \in G * \langle t \rangle$ satisfy $\text{ex}(w) = 1$. Then, after conjugation, w can be written as a product*

$$b_0 a_0^t b_1 a_1^t \cdots b_r a_r^t c t,$$

where $a_i \in Y, b_i \in X, i = 0, \dots, r$ and $c \in J$ for some $m > 0$.

Furthermore, provided w is not conjugate to gt for some $g \in G$, then $r \geq 0$ in the expression.

Remark The final sentence in the statement of lemma 2 is not given in [4], but is immediate from the proof given there.

Lemma 3 [4; Lemma 4.3, page 66] *Suppose that G is torsion-free, then any element a of Y is free relative to H . Similarly any element b of X is free relative to H' . \square*

We can now complete the proof of the main theorem, which follows closely the proof of [4; Theorem 4.4, pages 66–67]. By lemma 2 we can assume that $w = b_0 a_0^t b_1 a_1^t \cdots b_r a_r^t c t$, where $a_i \in Y, b_i \in X, i = 0, \dots, r$ and $c \in J$ and $r \geq 0$. We need to think of each a_i, b_i, c as functions of t and for clarity we shall introduce a new variable s . To be precise let

$$w(s, t) \equiv b_0(t) a_0^s(t) \cdots b_r(t) a_r^s(t) c(t) s$$

where s and t are independent variables.

Write Γ for $G * \langle t \rangle$ and let H, H' be the subgroups defined above. There is an isomorphism $\phi : H \rightarrow H'$ given by $h^\phi = h^t, h \in H$.

Lemma 3 gives the hypothesis of our key technical theorem in Section 4, which implies that

$$gs \text{ is never in the kernel of } \Gamma * \langle s \rangle \rightarrow \widehat{\Gamma}. \quad (**)$$

The case $m = 1$ is the special case (with t -shape $t^{-1} t t^{-1} \dots t t^{-1} t t$) covered by the proof in the last section, so we may assume that $m > 1$ and then $G \subset H \neq \emptyset$.

Each of the canonical elements of $a_i(t), b_i(t), c(t)$ is either in G or lies in H^t ; moreover in $\widehat{\Gamma}$ we have $h^s = h^\phi = h^t$ for each $h \in H$.

Since H is generated by elements of the form $t^{-i} g t^i$ for $i \leq m - 2$ and since $h^s = h^t$ for each $h \in H$ it follows by induction on i that we can freely exchange s and t in products of elements of the form $t^{-i} g t^i$ for $i \leq m - 1$. Thus we can exchange s and t in the coefficients of $w(s, t)$ and it follows that $w(s, s) = 1$ in $\widehat{\Gamma}$.

Now consider the following commutative diagram

$$\begin{array}{ccccc} \Gamma = G * \langle t \rangle & \subset & \Gamma * \langle s \rangle & \longrightarrow & \widehat{\Gamma} \\ \cup & & \cup & & \uparrow \\ G & \longrightarrow & G * \langle s \rangle & \longrightarrow & \widehat{G} \end{array}$$

where $\widehat{G} = \frac{G * \langle s \rangle}{\langle\langle w \rangle\rangle}$.

By (**), $gs \in \Gamma * \langle s \rangle$ does not map to $1 \in \widehat{\Gamma}$ for any $g \in G$. Therefore $gs \in G * \langle s \rangle$ never maps to $1 \in \widehat{G}$. This proves (*) as required. \square

5 The surjectivity problem and Whitehead torsion

If one adds n generators x_1, x_2, \dots, x_n and n relators w_1, w_2, \dots, w_n to a group G to form the group \widehat{G} then one can ask whether the natural homomorphism $G \longrightarrow \widehat{G}$ is injective. If it is injective then one can ask whether it is surjective. Our main theorem answers this question completely for torsion-free groups when $n = 1$.

The question of surjectivity, assuming injectivity, was raised by Cohen [2] in his study of Zeeman's conjecture. Assuming the natural homomorphism is injective then one can associate (after some normalization – see [9, pages 600-601]) to the set of words w_1, \dots, w_n a Whitehead torsion element $\tau \in \text{Wh}(G)$ (the Whitehead group of G). Cohen conjectured that if $\tau \neq 0$ then the injection cannot be onto. In closely related but independent work, in which he investigated inclusions of one 2-complex into another which are homotopy equivalences, Metzler [6] investigated the group theoretic combinatorics and the set of Whitehead torsion elements which are associated to such homotopy equivalences. He named the set of Whitehead torsion elements which can be realized by a relative 2-complex as $\text{Wh}^*(G)$

Our main theorem appears to give evidence that $\text{Wh}^*(G) = 0$. In fact, when $n = 1$ not only do we show that a necessary condition for surjectivity is that the torsion of the 1×1 matrix is 0, but we show that, up to homotopy of the attaching map, the added one- and two- cells can be collapsed away. However, one must be very cautious concerning what this means for $n > 1$ in that

- not all Whitehead torsion elements can be realized by 1×1 matrices, hence our result for $n = 1$ in no way answers the question of whether $\text{Wh}^*(G) = 0$,
- it is possible (an open conjecture) that $\text{Wh}(G) = 0$ for all torsion-free groups.

The only significant results on the surjectivity problem which we know of for $n > 1$ are those of Rothaus [9]. He develops an obstruction to the surjectivity of the map $G \rightarrow \widehat{G}$ in terms of representations of G into compact connected Lie groups. His theory had the following application for dihedral groups, whose Whitehead groups are known to be non-trivial.

Theorem [9; Theorem 11] *If $p \geq 5$ is an odd prime and $G = D_{2p}$ is the dihedral group of order $2p$ and n is any positive integer then there exist non-trivial Whitehead torsion elements such that every injective homomorphism $G \rightarrow \widehat{G} = \frac{G*\langle t_1, \dots, t_n \rangle}{w_1 \dots w_n}$ realizing this Whitehead torsion element is non-surjective.*

Beyond Rothaus' work, the surjectivity problem for $n > 1$ is an open and fascinating question.

6 An extension and a question

The main theorem (in the equivalent form given by lemma 1) can be extended:

Extension of Main Theorem

*Let G be a torsion free group and let $w \in G * \langle t \rangle$ be word whose t -shape is amenable (see [4, 5]) and consider the natural map $\pi: G \rightarrow \widehat{G} = \frac{G*\langle t \rangle}{\langle\langle w \rangle\rangle}$. Let $x \in G * \langle t \rangle$ be any word with t -shape t^n for some n .*

Then x is not in the kernel of π .

The proof is very similar to the proof of the main theorem. The cell e_∞^2 has n edges all oriented the same way (“uphill”). Notice that any other cell with an edge in common with e_∞^2 has its car traverse that edge in the “downhill” direction, since adjacent cells induce opposite orientations on a common edge. Choose any point $\omega \in \partial e_\infty^2$ not at a vertex. The flow constructed as in [4; page 68] for cells other than e_∞^2 has the property that there are times when all cars are going uphill and hence are not on ∂e_∞^2 . This leaves time for car A (on ∂e_∞^2) to rush round from just after ω to just before and hence there are no complete crashes on ∂e_∞^2 except at ω . This leads to the identical contradiction as in the proof of the main theorem.

The extension implies that all words of t -shape t^n for some n have infinite order in \widehat{G} . This leads to the natural question:

Question Suppose that G is torsion-free and that w is an amenable word. Is \widehat{G} torsion-free?

If the answer is yes, then we can deduce that $G \rightarrow \widehat{G}$ is never surjective when \widehat{G} is obtained from G by adding n generators and n relators *one pair at a time*.

References

- [1] **MM Cohen**, *A Course in Simple-Homotopy Theory*, GTM 10, Springer-Verlag, New York (1973)
- [2] **MM Cohen**, *Whitehead torsion, group extensions, and Zeeman's conjecture in high dimensions*, *Topology*, 16 (1977) 79–88
- [3] **Marshall M Cohen, Martin Lustig**, *The conjugacy problem for Dehn twist automorphisms of free groups*, *Comment. Math. Helv.* 74 (1999) 179–200
- [4] **Roger Fenn, Colin Rourke**, *Klyachko's methods and the solution of equations over torsion-free groups*, *L'Enseignement Mathématique*, 42 (1996) 49–74
- [5] **Roger Fenn, Colin Rourke**, *Characterisation of a class of equations with solutions over torsion-free groups*, from: “The Epstein Birthday Schrift”, (I Rivin, C Rourke and C Series, editors), *Geometry and Topology Monographs*, Volume 1 (1998) 163–17
- [6] **W Metzler**, *Über den Homotopietyp zweidimensionaler CW-Komplexe und Elementartransformationen bei Darstellungen von Gruppen durch Erzeugende und definierende Relationen*, *J. Reine Angew. Math.* 285 (1976) 7–23
- [7] **M Kervaire**, *On higher dimensional knots*, *Differential and combinatorial topology – a symposium in honour of Marston Morse*, *Princeton Math. Series*, 27 (1965)
- [8] **A Klyachko**, *Funny property of sphere and equations over groups*, *Comm. in Alg.* 21 (1993) 2555–2575
- [9] **OS Rothaus**, *On the non-triviality of some group extensions given by generators and relations*, *Ann. of Math.* 106 (1977) 599–612
- [10] **C P Rourke, B J Sanderson**, *Introduction to piecewise-linear topology*, Springer study edition, Springer-Verlag, Berlin (1982)